


The Architecture of Creation

Mathematical Structures, Generative Laws, and
the Recurrence of Form in Nature

Mathematical morphology · Physics · Biology · Complex systems



Contents

About This Report.....	5
Executive Summary	6
1. Central Thesis and Scope	8
1.1 The observation: form recurs	8
1.2 The thesis	9
1.3 What the thesis does and does not claim	9
1.4 Method and evidence standards	9
1.5 Intellectual context.....	10
2. Concept: What Is a Creation Structure?	11
2.1 Definition.....	11
2.2 Anatomy of a creation structure	11
2.3 The five-layer model.....	12
2.4 Law, constraint, mechanism and pattern.....	13
2.5 The possibility space	13
2.6 Intellectual lineage.....	14
2.7 Summary	14
3. Fundamental Structures: The Constraints Beneath Physical Reality.....	15
3.1 Why matter is repeatable	15
3.2 Symmetry and conservation	15
3.3 Energy minimisation and variational principles	15
3.4 Crystal symmetry and the limits of order	16
3.5 Scaling, dimensional analysis and universality	17
3.6 Summary of the fundamental layer	18
4. Geometric Structures: The Shapes the Constraints Favour.....	19
4.1 Spirals.....	19
4.2 Phyllotaxis: the pinecone problem.....	20
4.3 Hexagons and packing	23
4.4 Branching and transport networks.....	24
4.5 Self-similarity and fractals.....	25
4.6 Same geometry, same cause?.....	26
5. Biological Structures: How Living Systems Build Form.....	28
5.1 Genes set parameters, not shapes	28
5.2 Morphogens and positional information	28
5.3 Reaction–diffusion and Turing patterns.....	29
5.4 Cells as physical objects	30
5.5 Allometry and scaling	30
5.6 Body plans and morphospace	31
5.7 From gene to geometry: the pinecone revisited	32
6. Emergent Structures: Order Without a Designer.....	33
6.1 What emergence means here	33
6.2 Crystals and snowflakes.....	33
6.3 Patterns in fluids, chemistry and sand.....	34
6.4 Collective motion	34
6.5 Simple rules, complex outputs	35
6.6 Criticality and power laws	36
6.7 Type versus token	37
7. Reproductive and Evolutionary Structures: Why Nature Repeats Itself	38
7.1 Heredity as high-fidelity copying.....	38
7.2 Three routes to identical form.....	39
7.3 Deep homology	39
7.4 Convergent evolution	40
7.5 Adaptive landscapes and morphospace	41
7.6 Genotype–phenotype maps	41

7.7 Contingency and convergence	42
8. Synthesis: A Unified Framework	43
8.1 One architecture, five layers	43
8.2 Case studies across the layers	43
8.3 Interactions between layers	44
8.4 Why so much order?	45
9. Testing the Thesis: Predictions, Limits and Risks	46
9.1 Testable predictions	46
9.2 What would count against the thesis	46
9.3 Failure modes	47
9.4 Open questions	47
9.5 A note on interpretation	48
10. Applications and Heuristics	49
10.1 Where creation structures create value	49
10.2 Ten heuristics for reading a pattern	49
10.3 Research agenda	51
11. Conclusion	52
References	53
Primary literature	53
Books	55
Glossary	56
Appendix: Key Quantities	57

List of Figures

Figure 1. The five-layer creation-structure architecture used throughout this report	7
Figure 2. Recurrence across scales	8
Figure 3. Constraint filters nested inside the space of imaginable forms	12
Figure 4. Milestones in the study of generative form, 1917–2012	14
Figure 5. The crystallographic restriction	16
Figure 6. A Penrose tiling generated by subdividing golden triangles (Robinson construction, six	17
Figure 7. Metropolis simulation of the two-dimensional Ising model on a 128 by 128 lattice	18
Figure 8. Three spirals	19
Figure 9. Vogel's model with 900 seeds at four divergence angles	21
Figure 10. Size of the largest gap (99.5th percentile of the distance from a point in the pattern	21
Figure 11. Parastichies in a simulated seed head of 240 seeds placed at the golden angle	22
Figure 12. Perimeter required to enclose a cell of unit area	23
Figure 13. Voronoi cells for 120 random seeds (left) and after 60 rounds of relaxation towards	24
Figure 14. Murray's law	25
Figure 15. Recursive branching with different length ratios and angles	25
Figure 16. Iterated function systems	26
Figure 17. Positional information	29
Figure 18. Gray–Scott reaction–diffusion model on a 160 by 160 grid, simulated for 12,000 to 20,000	29
Figure 19. Alternative metabolic scaling exponents	31
Figure 20. Raup-style shell coiling	32
Figure 21. Snow-crystal growth in a Reiter-type cellular model on a hexagonal grid	34
Figure 22. A Vicsek-type simulation with 300 particles in a periodic box	35
Figure 23. Three elementary cellular automata	35
Figure 24. Avalanche sizes in a simulated Bak–Tang–Wiesenfeld sandpile on a 64 by 64 grid, after	36
Figure 25. Approximate error rates in DNA replication at successive levels of correction, and the	38
Figure 26. Convergent body form	40
Figure 27. An illustrative fitness landscape with three major peaks and two minor ones	41
Figure 28. A schematic genotype–phenotype map	42

Figure 29. Relative contribution of each layer to five recurring phenomena and to the case of.....44

Figure 30. A decision procedure for classifying the likely source of a recurring pattern 51

About This Report

This report develops and tests a single organising idea: that the recurrence of form in nature, from the spirals of a pinecone to the near-identity of unrelated animals that live the same way, is explained by *creation structures*, the rules, constraints and generative processes that determine which forms can emerge, how they develop and why they recur.

It is written for readers who want a rigorous but readable account: researchers scoping a programme, designers and engineers looking for transferable principles, and decision-makers who want to understand where the science is firm and where it is not.

How to read this report

Each section opens on a new page and can be read on its own. Section 1 states the thesis; Section 2 defines the concept and its five-layer structure; Sections 3 to 7 take the layers in turn; Section 8 synthesises them; Section 9 tests the thesis and lists its limits; Section 10 turns it into practical heuristics; Section 11 concludes.

Blue underlined text is a hyperlink. Links go to the primary literature where a specific result is cited (usually through its DOI or preprint record) and to a reference article where a concept is introduced. Every link was checked for resolution before publication. A full reference list appears at the end.

Evidence grades

Claims are graded where it matters, using a four-level scale.

Table 1. Evidence grades used in this report

Grade	Meaning
A: Established	Derived mathematically or confirmed repeatedly by independent experiment or observation.
B: Well supported	Strong evidence from several lines of work, with some unresolved detail or dissent.
C: Plausible or contested	Credible mechanism and partial evidence, but the field is actively debating it.
D: Speculative	A hypothesis offered for testing. It is labelled as such wherever it appears.

A note on the figures

All 30 figures were produced for this report. About half are original simulations or calculations of standard published models: phyllotaxis, Voronoi relaxation, reaction–diffusion, the Ising model, collective motion, cellular automata, a sandpile, snow-crystal growth and others. Each caption states what was computed. Figures marked *illustrative* are diagrams of a concept, not plots of measured data.

Executive Summary

The question

Pinecones, sunflowers and pineapples carry spirals whose counts are consecutive Fibonacci numbers. Honeycombs, basalt columns and graphene share a hexagonal geometry despite having nothing else in common. Sharks, ichthyosaurs and dolphins, separated by hundreds of millions of years of evolution, converged on the same streamlined body. Identical twins, cloned aspen groves and bacterial colonies are nearly indistinguishable. These recurrences are too frequent to be coincidence and too varied to share one cause. The question is what they have in common.

The thesis

Reality generates recurring forms because matter, energy, information and biological development operate within a constrained set of mathematical relationships and physical laws. The constraints define a small space of viable forms. Generative processes explore that space in a biased way, and selection and heredity filter and propagate what they find. Recurrence is the signature of a small possibility space explored by a biased generator.

We call the combination of constraints, generators and filters a *creation structure*. The thesis is pluralist: it does not claim that every pattern has one cause, but that every recurring pattern can be located within a layered architecture of causes.

Principal findings

1. **Recurrence has at least three distinct sources, and they should not be confused.** Shared physical constraint (hexagons in soap froth and basalt), shared developmental machinery inherited from a common ancestor (the Hox genes) and shared selection pressure acting on different lineages (the fusiform body) each produce similar forms for different reasons. Section 10 gives a decision procedure for telling them apart.
2. **Constraint is stronger than most people expect.** Symmetry, conservation laws, energy minimisation and geometry rule out most imaginable forms before biology begins. Only 2-, 3-, 4- and 6-fold rotational symmetries are compatible with a periodic lattice. Essentially one divergence angle packs growing seeds uniformly. Only a narrow band of branching ratios minimises transport cost.
3. **Simple local rules generate complex global order.** Reaction–diffusion systems, cellular automata and flocking models produce stripes, spots, fractals and coordinated motion without any central plan. The same small set of mechanisms recurs across chemistry, biology and physics.
4. **Biology uses physics as a collaborator, not an adversary.** Genes do not draw a pinecone. They set parameters (hormone transport, growth rate, cell adhesion) of processes whose geometry is then largely determined by physics. This is why many different genetic routes lead to the same shape.

5. **Evolution is both contingent and channelled.** Replaying life would not reproduce the same species, but it would revisit the same solutions to the same problems. The strongest evidence is the repeated independent origin of eyes, flight, C4 photosynthesis and streamlined bodies.
6. **The thesis is testable, and it can fail.** Section 9 lists eight predictions. The framework is weakened where recurrence turns out to be numerological coincidence, as with many claims about the golden ratio, or where historical accident, not constraint, explains similarity.

Implications

For science, creation structures offer a way to organise findings across physics, developmental biology and evolution. For engineering and design, they supply transferable generative rules (packing, branching, self-assembly) that can be exploited directly. For anyone reasoning about pattern, they provide a checklist: before asking *what does this pattern mean*, ask *which layer of the architecture produced it*. Section 10 condenses this into ten working heuristics.




Figure 1. The five-layer creation-structure architecture used throughout this report. Lower layers are more universal; higher layers carry more historical contingency.

1. Central Thesis and Scope

1.1 The observation: form recurs

Nature is not a catalogue of unrelated shapes. A small number of forms appear repeatedly, in systems that differ in material, scale and origin. Four examples frame the problem.

Spirals in seed heads. The scales of a pinecone, the florets of a sunflower and the leaves of a pineapple are arranged in two families of interlocking spirals. The number of spirals in each family is almost always a pair of consecutive terms of the [Fibonacci sequence](#): 5 and 8, 8 and 13, 21 and 34. The underlying arrangement, called [phyllotaxis](#), has been documented across many plant families that diverged long ago.

Hexagons in unrelated materials. A [honeybee comb](#), a sheet of [graphene](#), a field of cooled [basalt columns](#) and the cloud pattern at [Saturn's north pole](#) all display hexagonal order. The materials share no chemistry, and the length scales differ by more than seventeen orders of magnitude.

Near-identical animals. Sharks, tuna, ichthyosaurs and dolphins share a streamlined, fusiform body with a dorsal fin and a crescent-shaped tail. Ichthyosaurs (reptiles) and dolphins (mammals) descend from land-dwelling ancestors and re-evolved the shape independently. The phenomenon is called [convergent evolution](#). In other cases animals are identical for the opposite reason: they are [monozygotic twins](#) or members of a [clonal colony](#) and share a genome.

Branches everywhere. Rivers, lightning, lungs, neurons and trees all branch, and the diameters of their branches follow similar rules.




Figure 2. Recurrence across scales. Spirals, hexagons and branching networks appear in systems that differ in size by up to thirty orders of magnitude. Sizes are approximate characteristic lengths. Illustrative compilation by the author.

1.2 The thesis

The central thesis of this report is:

Reality generates recurring forms because matter, energy, information and biological development operate within a constrained set of mathematical relationships and physical laws. These constraints create a generative architecture in which different systems repeatedly converge on similar structures, proportions, geometries and organisational patterns.

The thesis rests on three linked claims.

1. **Constraint.** Mathematics and physics sharply restrict which forms are possible. Of all imaginable shapes, only a small fraction are stable, buildable, and consistent with conservation laws and geometry.
2. **Generation.** Processes that build form, from crystal growth to embryonic development, are not uniform samplers of the possible. They are biased by their own mechanics towards certain outcomes, so that a few forms are reached repeatedly from many starting points.
3. **Filtration.** Heredity propagates working forms and selection removes failing ones. When different lineages face the same problem, the same few solutions are repeatedly reached.

Together these claims explain recurrence as the output of a small possibility space explored by a biased generator and filtered by selection. Any system that satisfies the same constraints, runs the same kind of generator and faces the same filter will tend to produce the same form.

1.3 What the thesis does and does not claim

The thesis is deliberately modest in some respects, and strong in others.

It does not claim a single cause. Honeycombs, snowflakes and eyes arise through different mechanisms. The claim is that each can be placed within the same architecture, not that one law explains all of them.

It does not claim design, purpose or teleology. Recurrence arising from constraint and selection requires no foresight. This report takes no position on metaphysical interpretations; Section 9.5 discusses why the question is raised and why the science does not depend on it.

It does not endorse numerology. Many popular claims about the [golden ratio](#) in art, architecture and the human body do not survive scrutiny. The thesis is aimed at patterns that arise from identified mechanisms and can be quantified, tested and predicted.

It does claim predictive content. If recurrence reflects a small possibility space, then (a) independently evolved solutions to the same problem should cluster, (b) simulating the generative rules should reproduce the forms without a blueprint, and (c) perturbing the constraints should shift the forms in predictable ways. Section 9 sets out specific tests.

1.4 Method and evidence standards

The report is a synthesis of established results across mathematics, physics, developmental biology and evolutionary theory, supplemented by original simulations. Each claim is graded on the A to D scale defined at the front of this report.

Three methodological commitments shape the analysis.

- **Mechanism before metaphor.** A recurring pattern is only explained when a mechanism is identified that generates it and can be modelled. Resemblance alone is not evidence.
- **Quantification where possible.** Wherever the literature permits, patterns are described by measured quantities (angles, exponents, error rates) not by visual impression.
- **Null models.** For each pattern we ask what random or unconstrained processes would produce, since recurrence is only surprising relative to a null expectation. This guards against the [Texas sharpshooter fallacy](#) and [pareidolia](#).

1.5 Intellectual context

The idea that form follows from rule has a long lineage. [D'Arcy Thompson's](#) 1917 book [On Growth and Form](#) argued that biological form could be understood through physical forces and mathematical transformation. [Alan Turing's](#) 1952 paper showed that interacting chemicals could spontaneously produce spots and stripes. The modern programme draws on [dissipative structures](#), [fractal geometry](#), [evolutionary developmental biology](#) and [complex systems theory](#). Section 2.6 presents the lineage as a timeline.

The aim here is to organise that work under one structure and to test where the organising idea holds and where it fails.

2. Concept: What Is a Creation Structure?

2.1 Definition

A **creation structure** is the set of underlying rules, constraints and generative processes that determine which forms can emerge, how they develop, and why particular patterns recur across different systems.

The definition has four parts, each of which matters.

- **Rules and constraints** specify what is *allowed*. They are the mathematical and physical limits on form.
- **Generative processes** specify what is *produced*. They are the mechanisms (growth, diffusion, self-assembly, development) that turn constraints and inputs into shapes.
- **Development** specifies *how* a form arises over time, from simple initial conditions to a finished structure.
- **Recurrence** is the observable consequence: the same output from different systems.

A creation structure is therefore not a shape, but the machinery behind a family of shapes. A pinecone is an instance. The *creation structure* is the combination of seed-packing geometry, hormone-driven primordium placement and growth physics that produces pinecones and sunflowers alike.

2.2 Anatomy of a creation structure

Every creation structure we examine has the same five components.

Table 2. Components of a creation structure

Component	Role	Example (phyllotaxis)
Constraint	Defines the space of allowed forms.	Primordia cannot overlap; growth proceeds from a central meristem.
Generator	The process that produces form over time.	Primordia form at the point of least inhibition around the growing tip.
Medium	The material and energy supply in which the process runs.	Plant tissue; auxin transport; cell growth.
Selector	A filter that keeps or removes outcomes.	Natural selection favours seed packing that fills space without gaps.
Attractor	A stable outcome reached from many starting points.	The golden-angle divergence and Fibonacci parastichy counts.

The *attractor* is the concept that most directly explains recurrence. In dynamical systems an [attractor](#) is a state towards which a system evolves from a wide range of initial conditions. When a generator has an attractor, many different histories converge on the same form, and the form is *robust*: it is


reached again and again, despite noise. This is why a pinecone's spirals do not depend on the exact conditions of each tree.

2.3 The five-layer model

The report organises creation structures into five layers, following the structure of the question. Each layer addresses a different part of the problem, uses different tools, and carries a different degree of universality and historical contingency.

Table 3. The five layers of creation structure

Layer	Central question	Tools	Typical output
1. Fundamental	What configurations are possible at all?	Mathematics, symmetry, conservation, variational principles	Crystal symmetries, minimal surfaces, scaling laws
2. Geometric	Which shapes do the constraints favour?	Geometry, packing theory, fractals, network theory	Spirals, hexagons, branching, tilings
3. Biological	How do living systems build form?	Developmental genetics, morphogens, cell mechanics	Body plans, stripes, organ shape
4. Emergent	How does order arise without a plan?	Dynamical systems, statistical physics, simulation	Snowflakes, flocks, convection cells
5. Reproductive and evolutionary	Why are the same forms found again and again?	Population genetics, phylogenetics, selection theory	Homology, convergence, clones



Each layer is a constraint filter. The innermost region is small relative to what is imaginable.

Figure 3. Constraint filters nested inside the space of imaginable forms. Each layer removes most candidates; the realised set is small. Illustrative.

Two properties of this ordering are important.

Universality decreases upward. The conservation of energy applies to a snowflake and an elephant alike. The Hox gene complex applies only to animals. The genome of a particular species applies to that species. Lower layers explain *wide* recurrence; higher layers explain *narrow* recurrence.

Contingency increases upward. Physical law has no history. Development has an evolutionary history that fixes some of its details. Evolution depends on specific events (mutations, extinctions, geography). The layers therefore also order the degree of *historical accident* we should expect in an explanation.

2.4 Law, constraint, mechanism and pattern

Much confusion about recurrence comes from running together four different kinds of statement.

Table 4. Four kinds of statement, often confused

Term	What it is	Example
Law	A regularity that holds universally under stated conditions.	Conservation of energy; the second law of thermodynamics.
Constraint	A restriction on what a system can do, arising from laws plus the system's structure.	A periodic crystal cannot have five-fold rotational symmetry.
Mechanism	A causal process that produces an outcome.	Short-range activation with long-range inhibition produces spots.
Pattern	The observed regularity in the output.	Leopard spots; Fibonacci spiral counts.

A law is not a pattern. A pattern is only *explained* when a mechanism is exhibited, operating under constraints that follow from laws. When we say a pinecone's spirals are "caused by the Fibonacci sequence" we are making a category error: the Fibonacci numbers describe the output, and the cause is a growth and packing process (Section 4.2).

2.5 The possibility space

A useful way to express the thesis is geometrically. Imagine the set of all conceivable forms. Mathematical consistency removes some. Physical law removes more. Developmental accessibility, meaning whether a growth process can reach a form from a feasible starting point, removes still more. Finally, selection and heredity determine which of the accessible forms persist.

At each stage the set shrinks, and what remains is far smaller than the starting space. Observed forms occupy a tiny region. Recurrence follows because different systems, exploring the same shrunk space with the same biased generators, often land in the same small neighbourhoods.

This picture has a testable implication. If the possibility space really is small, the forms that biology has *not* produced should include many that physics permits but development cannot reach, and biologists should be able to identify such forbidden-but-possible forms (Section 5.6 discusses shell morphospace, where this has been done quantitatively).

2.6 Intellectual lineage



Figure 4. Milestones in the study of generative form, 1917–2012. The timeline is selective, and each entry corresponds to a result discussed in this report.

Three threads run through this history. The first is *geometric*: Thompson, Mandelbrot and Hales sought the mathematical regularities in form. The second is *dynamical*: Turing, Prigogine and the complexity theorists showed that order can emerge from local interactions. The third is *evolutionary and developmental*: Wolpert, Gould and Lewontin, and the founders of evo-devo asked how a shared body of rules is inherited and tuned. The creation-structure idea is an attempt to treat the three threads as parts of one architecture.

2.7 Summary

Key points. A creation structure is a layered combination of constraints, generators and filters. Recurrence is explained when a mechanism, operating within constraints, is shown to reach the same attractor from many starting points. The five-layer model orders the explanations from universal (physics) to contingent (history). A law is not a mechanism, and a mechanism is not a pattern: keeping them apart prevents most errors of interpretation.

3. Fundamental Structures: The Constraints Beneath Physical Reality

Layer 1. Mathematical relationships, physical laws, symmetry, conservation principles and the properties of matter and energy determine which configurations are possible.

3.1 Why matter is repeatable

The deepest reason that forms recur is that the building blocks are *identical*. Every electron has exactly the same mass, charge and spin as every other. In quantum mechanics this is stronger than similarity: particles of the same kind are [indistinguishable](#), and swapping two of them changes nothing observable. The consequences are physical and large. The [Pauli exclusion principle](#), which follows from the exchange symmetry of electrons, forces atoms into shell structures, and the shell structures produce the [periodic table](#). Every atom of carbon-12 in a pinecone and in a diamond is the same object.

This matters for the thesis because it makes form *reproducible*. If matter came in unlimited unique varieties, recurrence would be impossible. Because it comes in a few dozen stable kinds with fixed properties, the same chemistry, the same bond angles and the same crystal lattices appear whenever the conditions repeat. Evidence grade: **A**.

3.2 Symmetry and conservation

In 1918 [Emmy Noether](#) proved that every continuous symmetry of a physical system corresponds to a conserved quantity. Symmetry under shifts in time gives conservation of energy; under shifts in space, conservation of momentum; under rotations, conservation of angular momentum. Conservation laws are therefore not independent facts but consequences of symmetry, and they sharply restrict what a system can do.

Symmetry also explains why patterns appear. When the laws governing a system are symmetric but the system's state is not, [symmetry breaking](#) has occurred. A pencil balanced on its tip obeys rotationally symmetric laws, yet it falls in a particular direction. A ferromagnet cooled below its critical temperature picks one magnetic orientation. A fertilised egg, spherically symmetric to a first approximation, develops a head and a tail. In each case the *form* that results is limited by the symmetry of the underlying laws, because the broken states are related to one another by the original symmetry.

3.3 Energy minimisation and variational principles

A large class of physical systems settle into the state of lowest energy, or follow the path that makes an *action* stationary (the [principle of least action](#)). Many apparently different phenomena share this logic.

A [soap film](#) spanning a wire frame adopts the surface of least area, because surface tension penalises area. The empirical rules, known as [Plateau's laws](#), follow: films meet three at a time along an edge at exactly 120 degrees, and four edges meet at a vertex at the tetrahedral angle of about 109.5 degrees. No other junctions are stable. The same rules govern the arrangement of bubbles in

foam, the boundaries of cells in many tissues and the cracks in some cooled materials. The geometry is imposed by minimisation, not designed.

The consequence for the thesis is direct. Where a system minimises an energy or a cost, the *family* of forms it can adopt is narrowed to the solutions of an optimisation problem. If different systems face the same optimisation problem, they converge on the same solution regardless of material.

3.4 Crystal symmetry and the limits of order

Crystals are built by repeating a unit in space. The mathematical structure of such repetition is completely classified: there are 17 [wallpaper groups](#) in two dimensions, 32 crystallographic [point groups](#) and 230 [space groups](#) in three. The classification is complete, so every periodic crystal ever discovered belongs to one of these classes.

A key result is the [crystallographic restriction theorem](#): a periodic lattice can have only 2-, 3-, 4- or 6-fold rotational symmetry. Five-fold, seven-fold and higher symmetries are forbidden. The reason is geometric. Three regular pentagons meeting at a point leave a gap of 36 degrees, because each pentagon's interior angle is 108 degrees and three of them cover only 324 of the available 360. No arrangement of pentagons fills the plane without gaps or overlaps.




Figure 5. The crystallographic restriction. Triangles, squares and hexagons tile the plane without gaps; regular pentagons leave a 36 degree gap at each vertex and cannot. Original diagram.

This explains a familiar fact, that snow crystals have six arms, never five. Water molecules in ice Ih sit on a hexagonal lattice, and the macroscopic crystal inherits that symmetry (Section 6.2).

The restriction applies to *periodic* order only. In 1984 Dan Shechtman and colleagues reported an [aluminium–manganese alloy](#) whose diffraction pattern showed sharp spots with ten-fold symmetry. It was ordered but not periodic: the first [quasicrystal](#). The discovery, recognised by the 2011 Nobel Prize in Chemistry, shows that the boundary of the possible is itself a mathematical question, not a fixed barrier. The mathematical model is the [Penrose tiling](#), which fills the plane with two tile shapes in a pattern that has five-fold symmetry and never repeats. The ratio of the numbers of the two tile types is the golden ratio.




Figure 6. A Penrose tiling generated by subdividing golden triangles (Robinson construction, six subdivisions). The pattern has local five-fold symmetry but no translational period. Original calculation.

Evidence grade: **A** for the classification and the restriction theorem; **A** for the existence of quasicrystals.

3.5 Scaling, dimensional analysis and universality

Some constraints come from nothing more than arithmetic. If an object is scaled up by a factor L , its surface area grows as L^2 and its volume as L^3 . This [square–cube law](#), noted by Galileo, limits the size of any structure supported by strength per area and explains why an insect scaled to the size of a horse would collapse and a whale cannot live on land. [Dimensional analysis](#) generalises the idea: a physical relationship can only combine quantities in ways that keep units consistent, which often fixes its form up to a constant.

A more profound constraint is *universality*. Near a continuous [phase transition](#), systems that differ in all microscopic detail show the same large-scale behaviour, described by the same *critical exponents*. A liquid–gas critical point and a uniaxial magnet belong to the same universality class. The explanation, developed by Kenneth Wilson in the [renormalisation group](#), is that microscopic details are progressively averaged away on larger scales; only the dimension of space and the symmetry of the order parameter survive.

The [Ising model](#), a lattice of spins that each point up or down and prefer to align with neighbours, is the simplest example. Its exact solution in two dimensions gives a critical temperature of about 2.269

in units of the coupling. Below it the lattice orders; above it the lattice is random; at the critical point the lattice shows structure on every scale.




Figure 7. Metropolis simulation of the two-dimensional Ising model on a 128 by 128 lattice. Left: below the critical temperature the lattice is ordered. Centre: at the critical temperature clusters appear on all scales. Right: above it the lattice is disordered. Original simulation.

Universality is the clearest example of a creation structure at the fundamental layer: it says that *many different microscopic realisations converge on one large-scale form*. This is what the thesis claims at all layers, and here it is proven. Evidence grade: **A**.

3.6 Summary of the fundamental layer

Table 5. Fundamental structures and what they constrain

Structure	Statement	Constrains	Grade
Identical particles	Particles of one kind are indistinguishable.	Chemistry, the periodic table	A
Noether's theorem	Continuous symmetries imply conserved quantities.	Allowed dynamics	A
Variational principles	Systems settle where energy or action is stationary.	Soap films, foams, cell boundaries	A
Crystallographic restriction	Periodic lattices allow only 2-, 3-, 4-, 6-fold rotations.	Crystal and snowflake symmetry	A
Square–cube law	Area scales as L^2 , volume as L^3 .	Size limits and shape of organisms	A
Universality	Microscopic detail is lost near criticality.	Phase transitions, scaling	A

Key points. The fundamental layer fixes the menu. It guarantees that the same ingredients behave the same way everywhere, restricts structures to those compatible with symmetry and conservation, and makes different systems converge when they solve the same optimisation problem. It does not say which forms nature *will* produce, only which it cannot.

4. Geometric Structures: The Shapes the Constraints Favour

Layer 2. Spirals, branching networks, hexagonal arrangements, radial symmetry, fractal-like structures and repeated proportions arise under particular combinations of growth, packing and physical forces.

4.1 Spirals

Spirals fall into three mathematically distinct families. In an Archimedean spiral the radius grows linearly with angle, as in a coiled rope. In a [logarithmic spiral](#) the radius grows exponentially with angle, so the curve crosses every ray from the centre at the same angle, and enlarging it by any factor gives the same curve rotated. The [golden spiral](#) is a logarithmic spiral whose radius grows by the golden ratio every quarter turn.




Figure 8. Three spirals. Left: Archimedean. Centre: logarithmic, with growth factor equal to the golden ratio per quarter turn. Right: quarter-circle arcs inscribed in Fibonacci squares, an approximation to the golden spiral. Original construction.

The logarithmic spiral is the natural shape of anything that grows at its open end while keeping its form. A [mollusc shell](#) adds material at the aperture; each new increment is a scaled copy of the previous one, and the shell winds around as one side grows slightly faster than the other. The shape is preserved as size increases, which is what a logarithmic spiral means. This argument, due to D'Arcy Thompson, explains the spiral in snail shells, ram's horns, elephant tusks and claws.

A caution is needed. It is often stated that the nautilus shell follows the golden spiral. It does not. The nautilus shell widens by a factor of roughly three per whorl in standard measurements, while a golden spiral widens by about 6.85 per whorl. The nautilus is a logarithmic spiral, and the golden spiral is one particular member of that family. Likewise, the spiral arms of a hurricane or a galaxy are spirals by different mechanisms (convergent winds and the Coriolis effect; density waves in a rotating disc), so geometric similarity does not imply common cause. Evidence grade: **A** for the geometry and the growth argument; the golden-ratio claim for shells is **false**.

4.2 Phyllotaxis: the pinecone problem

Why do pinecones, sunflowers and pineapples show Fibonacci spiral counts? The [phyllotaxis](#) literature has a complete answer, and it illustrates the whole creation-structure framework in miniature.

The observation. Successive leaves or seeds are placed around a central axis at a nearly constant *divergence angle*, which in most Fibonacci-type plants is close to 137.5 degrees, the [golden angle](#). It equals 360 degrees divided by the square of the golden ratio. In a sunflower head, counting the visible spirals in each direction gives pairs such as 34 and 55 or 55 and 89; in a pinecone, 5 and 8 or 8 and 13.

The geometry. In 1979 Helmut Vogel proposed a simple model: place the n -th seed at angle n times the divergence angle and at distance proportional to the square root of n . The result depends sharply on the angle. If the angle is a rational fraction of a full turn, seeds line up on rays and leave wasted gaps. If it is close to a simple fraction, the seeds form a few wide spirals with gaps between them. Only when the angle is badly approximated by every rational number does the packing become uniform, and the angle that is worst approximable by rationals is the golden angle, whose continued fraction expansion consists entirely of ones.




Figure 9. Vogel's model with 900 seeds at four divergence angles. Only the golden angle (top left) fills the disc uniformly. A deviation of half a degree (top right) produces visible gaps, and simple rational angles (bottom) collapse the pattern onto rays. Original simulation.




Figure 10. Size of the largest gap (99.5th percentile of the distance from a point in the pattern to the nearest seed) for an 800-seed head as the divergence angle varies. Rational angles produce spikes of large gaps. The golden angle sits among the lowest values (about 1.1, against 8.1 at 135 degrees and 5.6 at 150 degrees). Original calculation.

The spiral counts follow. Once seeds are placed at the golden angle, the neighbours of each seed fall at index differences that are Fibonacci numbers. The visible spirals (parastichies) connect seeds whose index differs by a Fibonacci number, so the counts of the two spiral families are consecutive Fibonacci numbers. The Fibonacci counts are a *consequence* of the angle, not a separate fact.

8 spirals

13 spirals

Consecutive Fibonacci numbers
(8, 13) — as counted on pinecones




Figure 11. Parastichies in a simulated seed head of 240 seeds placed at the golden angle. Joining seeds whose index differs by 8 (blue) and by 13 (amber) traces the two interlocking spiral families seen on pinecones. Original simulation.

The dynamics. Why do plants use the golden angle at all? In 1992 Stéphane Douady and Yves Couder showed in a [physical experiment](#) that no plant is needed. Droplets of magnetised fluid fell at regular intervals onto the centre of a dish in a magnetic field, and repelled one another as they moved outwards. The droplets arranged themselves in Fibonacci spirals, and the divergence angle approached the golden angle as the control parameter (the ratio of the time between droplets to their travel time) decreased. The mechanism is *each new element appears where the inhibition from the existing ones is smallest*. The golden angle is the attractor of this dynamic. The pattern is robust because many different parameter values lead to it.

The molecules. In plants the inhibition is implemented by the plant hormone auxin. [Reinhardt and colleagues](#) showed in 2003 that the sites at which new organs form at the shoot tip are set by auxin accumulation, which in turn is controlled by the polar transport protein PIN1. Existing primordia drain auxin from their surroundings, creating an inhibition zone; the next organ appears where auxin concentration first rises above threshold. The molecular machinery realises the geometric rule.

Caveats. Not every plant is Fibonacci. Alternative sequences (Lucas numbers, doubled Fibonacci) and non-spiral arrangements (opposite, whorled) occur and are better explained by the same dynamics with different parameters, in particular the relative size of the primordia and the growth rate of the meristem. A minority of species also show irregular or changing patterns. This is consistent with the thesis, because the framework predicts a small family of attractors, selected by parameters, not one forced outcome.

Evidence grade: **A** for the geometry of packing; **B** for the dynamical model; **B** for the molecular implementation.

4.3 Hexagons and packing

Hexagons recur because they solve two problems at once: covering a surface without gaps, and doing so with the least boundary.

The honeycomb conjecture. Among all ways of dividing a plane into regions of equal area, a regular hexagonal grid has the least total perimeter. The statement was believed since antiquity and proved by Thomas Hales in 1999, published in [Discrete and Computational Geometry](#) in 2001. For a cell of unit area, a triangle needs a perimeter of 4.56 and a square of 4.00, but a hexagon only 3.72. The saving is about 7% relative to squares. The only way to be cheaper is a circle, which does not tile.




Figure 12. Perimeter required to enclose a cell of unit area. Hexagons are the most economical shape that tiles the plane. Original calculation.

Bees may not be geometers. It is tempting to credit bees with solving an optimisation problem. [Karihaloo and colleagues](#) proposed in 2013 that bees build circular cells and that surface tension acts on the warm, softened wax to pull them into rounded hexagons where they meet. Whether the hexagons are built or flow into shape is still debated, and the debate shows the point: the geometry can arise through physics at the level of the material, without requiring a built-in plan. Evidence grade: **B** (the optimality proof) and **C** (the mechanism of comb formation).

A topological necessity. For any planar network in which three cells meet at each vertex, [Euler's formula](#) forces the average number of sides per cell to be six (exactly on a torus, and asymptotically in a large planar patch). Real tissues, foams and cracks have many five- and seven-sided cells, but the mean is pinned at six. In 2006 [Gibson and colleagues](#) found that proliferating epithelia in fruit-fly, frog and Hydra tissues share a characteristic distribution of cell shapes in which hexagons are the

most common class and the average is six sides, and they traced the distribution to the geometry of cell division. A constraint from topology and a growth process work together.

Voronoi cells. If space is divided so that each point belongs to the nearest of a set of seeds, the cells form a [Voronoi diagram](#). Many natural cell patterns resemble Voronoi diagrams, because they arise by growth outward from seeds at similar rates. If the seeds are random, the cells are irregular. If the seeds are evenly spaced, as produced by [Lloyd's algorithm](#) or by mutual repulsion, the cells approach hexagons.




Figure 13. Voronoi cells for 120 random seeds (left) and after 60 rounds of relaxation towards evenly spaced seeds (centre). Teal: six sides; blue: five; amber: seven; pink: eight or more; grey: four. Right: among interior cells, 33% were hexagonal for random seeds and 80% after relaxation. Original simulation.

Packing in three dimensions. The corresponding result for spheres is the [Kepler conjecture](#): the densest packing of equal spheres, filling about 74% of space, is the face-centred cubic arrangement used by grocers for oranges. It was proved by Hales and collaborators, with a [formal computer-checked proof](#) published in 2017. Close-packed arrangements underlie most metal crystal structures and the layering of many biological assemblies.

4.4 Branching and transport networks

A branching network solves a different problem: delivering a flow from a source to many destinations, or collecting from many sources, at the lowest cost.

Murray's law. In 1926 Cecil Murray reasoned that a blood vessel has two costs: the power needed to pump fluid through it, which falls steeply as the vessel widens, and the metabolic cost of maintaining the blood and the vessel wall, which rises with volume. Minimising the sum gives flow proportional to the cube of the radius, and [Murray's law](#) at a branch point: the cube of the parent radius equals the sum of the cubes of the daughter radii. For a symmetric split into two equal branches, each daughter has 0.79 of the parent's diameter. Vascular, respiratory and plant transport networks approximately follow the rule, with deviations in specific cases that are informative about additional constraints. Evidence grade: **B**.




Figure 14. Murray's law. Left: at a symmetric bifurcation each daughter has 0.79 of the parent diameter. Right: over many generations this taper is far gentler than halving of the diameter at every split. Original calculation.

One rule, many trees. Recursive branching, where each branch ends in two smaller copies of itself, generates a wide range of natural-looking forms from two parameters: the length ratio of successive generations and the branching angle. Systems of this type are called [L-systems](#) after Aristid Lindenmayer, who introduced them to model plant growth.




Figure 15. Recursive branching with different length ratios and angles. Nine generations in each case. Original construction.

Branching by growth instability. Rivers, lightning and some mineral dendrites do not branch to minimise cost. They branch because a growing tip in a diffusing field becomes unstable and sprouts side branches, a process modelled by [diffusion-limited aggregation](#). The visual resemblance to a lung or a tree conceals a different cause, so branching is a pattern with at least two creation structures behind it: *optimisation* (vessels, trees) and *instability* (rivers, lightning, dendrites). Evidence grade: **B**.

4.5 Self-similarity and fractals

[Benoit Mandelbrot](#) showed in 1967 that the length of a coastline is not a fixed number but depends on the ruler: shorter rulers resolve more bays and measure a longer coast. The length grows as a power of the inverse ruler size, with an exponent that defines a fractal dimension, about 1.25 for the west coast of Britain. This property, in which structure recurs at every scale, is called [self-similarity](#).

The generative content is clearest in [iterated function systems](#), where a handful of simple transformations applied repeatedly produce a complex limit shape. Three half-scale contractions

towards the corners of a triangle generate the [Sierpiński triangle](#); four affine maps generate the [Barnsley fern](#), a leaf-shaped fractal.




Figure 16. Iterated function systems. Left: three maps produce the Sierpiński triangle. Right: four affine maps (with the probabilities 1, 85, 7 and 7 per cent) produce a fern-like shape. Each panel shows more than 100,000 points. Original simulation.

This is a measurable form of the creation-structure idea: a short rule set (here, a few dozen numbers) specifies a form whose visible detail is vastly larger. The *compressibility* of natural form is evidence that a compact generator is at work.

A necessary caution: real objects are fractal only over a limited range of scales, usually between one and two decades. [Avnir and colleagues](#) found in 1998 that many published claims of fractal scaling in nature rested on very narrow ranges. Fractal geometry is a useful description where it holds over a wide range, and a pretence where it does not. Evidence grade: **A** (mathematics) and **B** (natural occurrences).

4.6 Same geometry, same cause?

Recurrence of geometry across scales does not by itself show a common cause. Three situations must be distinguished.

Table 6. When similar geometry does and does not indicate a common cause

Pattern	Systems	Mechanism	Common cause?
Hexagonal cells	Foams, epithelia, basalt, comb	Energy minimisation and topology (shared constraint)	Yes, at the level of constraint
Fibonacci phyllotaxis	Sunflower, cone, pineapple	Inhibitory field dynamics (shared attractor)	Yes, within plants
Logarithmic spirals	Shells, horns, hurricanes, galaxies	Growth that preserves shape vs. fluid dynamics vs. density waves	Only within shells and horns
Branching	Vessels, trees; rivers, lightning	Cost minimisation vs. growth instability	No: two classes

Key points. The geometric layer shows how a few simple rules generate recurring shapes. The same Voronoi or energy-minimisation logic reappears in unrelated materials, and a single dynamic (inhibition from existing elements) explains both the golden angle and Fibonacci spiral counts in plants. But resemblance is not identity: spirals and branches have several unrelated causes, and each claim of commonality must be verified by identifying the mechanism.

5. Biological Structures: How Living Systems Build Form

Layer 3. Genetic regulation, developmental pathways, cell behaviour, natural selection and physical constraints generate recognisable similarities within and across species.

5.1 Genes set parameters, not shapes

A common picture treats the genome as a blueprint, with a gene for each feature. Developmental biology shows a different picture. Genes encode proteins that act as signals, receptors, adhesion molecules and switches. The embryo uses them to run processes (growth, signalling, cell division) whose *geometric outcome* is determined jointly by the genes and by physics. A gene sets a parameter; the form is the output of the process.

The strongest evidence is how much of the machinery is shared. The [Hox genes](#), which specify position along the head-to-tail axis, were discovered through mutations in fruit flies that converted one body segment into another. The same gene family, in the same order along the chromosome, patterns the body axis in mice and humans. [Pax6](#), a master regulator of eye development, can trigger ectopic eyes on the legs and wings of flies when expressed there, and the mouse version of the gene works in the fly. Sean Carroll and colleagues summarised the broader pattern in the idea of a [developmental toolkit](#): a small, ancient set of regulatory genes used again and again to build very different animals.

[Davidson and Erwin](#) argued in 2006 that the toolkit is organised into gene regulatory networks whose core "kernels" are deeply conserved and difficult to change, because altering them damages many downstream processes at once. This produces *developmental constraint*, a bias in which forms can be produced. Maynard Smith and colleagues [defined it](#) in 1985 as a bias in the production of phenotypic variants caused by the structure and dynamics of the developmental system. Evidence grade: **A** for the conservation of the toolkit; **B** for the extent to which it constrains evolution.

5.2 Morphogens and positional information

How does an embryo, made of cells with identical genomes, assign different fates at different places? In 1969 Lewis Wolpert proposed that cells read their position from a concentration gradient and compare it with thresholds, a scheme he illustrated with the [French flag](#) problem: if a row of cells must become blue, white and red in fixed proportions, a gradient with two thresholds is sufficient. The diffusible signals are called [morphogens](#).

The model has direct support. In the fruit-fly embryo the protein [Bicoid](#) forms a concentration gradient from the head end, and cells activate different genes at different concentrations. A morphogen that is produced at one end, diffuses and is degraded generates an exponentially decaying profile whose length scale is set by the ratio of diffusion rate to degradation rate. The thresholds then partition the tissue into domains.




Figure 17. Positional information. A morphogen gradient with two thresholds divides a tissue into three fates (A, B, C). If the gradient's length scale adjusts to the size of the tissue, as in the right panel, the same proportions are achieved in a smaller embryo. The curves are exponentials; the figure is illustrative.

The scheme has a built-in difficulty that makes it a good example of a constraint: embryos of different sizes produce animals with the same proportions, so the gradient must scale with tissue size. Several mechanisms for achieving that scaling have been proposed and tested, and the question remains an active field. What is not in doubt is the *design logic*: a small set of parameters (source strength, diffusion, decay, thresholds) generates reproducible proportions. Evidence grade: **A** for the gradient principle; **B** for its universal application.

5.3 Reaction–diffusion and Turing patterns

In 1952 Alan Turing asked how a uniform tissue could break symmetry and form a pattern. He showed that two interacting chemicals that diffuse at different rates could be unstable to small perturbations and spontaneously produce periodic spots or stripes. The essential ingredients, clarified by [Gierer and Meinhardt](#) in 1972, are short-range activation and long-range inhibition: a substance that promotes its own production and also produces a faster-spreading inhibitor.

Reaction–diffusion systems are one of the clearest examples of a creation structure. A single pair of equations produces spots, stripes or labyrinths depending on a few parameters. The Gray–Scott model, whose parameter space was mapped by [John Pearson](#) in 1993, shows this directly.




Figure 18. Gray–Scott reaction–diffusion model on a 160 by 160 grid, simulated for 12,000 to 20,000 steps. The same two equations produce spots, stripes and a labyrinth as the feed rate F and kill rate k change. Original simulation.

The question that matters for the thesis is whether living tissues use this mechanism. The evidence has accumulated over three decades.

- In the marine angelfish *Pomacanthus*, [stripes rearrange continuously](#) as the fish grows, in the way a Turing pattern should adjust, rather than the way a fixed prepattern would.
- In mice, the spacing of [hair follicles](#) depends on a Wnt activator and a Dkk inhibitor that behave as a reaction–diffusion pair.
- In the developing limb, digit spacing is controlled by a [Bmp–Sox9–Wnt network](#). [Sheth and colleagues](#) showed that reducing the dose of Hox genes increases the number of digits and makes them thinner, as expected if Hox genes tune the wavelength of a Turing-type mechanism.
- In zebrafish, stripes arise from interactions between pigment cells with short-range and long-range effects of opposite sign, a [cellular analogue](#) of a Turing system.

A further prediction of reaction–diffusion theory is geometric. In narrow domains, spots are not stable and stripes must form. J. D. Murray pointed out that this predicts that an animal can have a spotted body and a striped tail, but not a striped body and a spotted tail. The rule fits cheetahs, leopards and genets. Evidence grade: **B** (strong for digits, hair follicles and fish; model-dependent for many other cases). A useful review is by [Kondo and Miura](#).

5.4 Cells as physical objects

Tissues are built from cells that adhere, push, pull and divide, and the forces involved obey mechanics. Two examples show how this reduces the number of free parameters.

Differential adhesion. In 1963 Malcolm Steinberg proposed that cells sort themselves by *adhesion strength*, in the same way that oil and water separate. If two cell types are mixed and one is more cohesive, the cohesive type forms a ball in the middle and the other envelops it. The [differential adhesion hypothesis](#) predicts that tissues behave as liquids with surface tension and settle into the arrangement of lowest adhesive energy. Layered tissues, organ boundaries and the arrangement of the germ layers show features consistent with this.

Buckling and folding. Folded structures arise when a growing layer is constrained by a substrate. [Tallinen and colleagues](#) modelled the human brain's cortical folds by swelling a gel layer on a gel core shaped like a fetal brain, and the resulting pattern of ridges and grooves resembled the real one. If a physical model without any genetic instructions reproduces the typical folding pattern, a large part of the fold geometry is explained by mechanics. Evidence grade: **B**.

5.5 Allometry and scaling

Biological form changes with size in regular ways. The study of this scaling is called [allometry](#). The best-known result is [Kleiber's law](#): across animals from mice to elephants, resting metabolic rate scales with body mass raised to a power close to three-quarters, not to the two-thirds that simple surface-area arguments predict.




Figure 19. Alternative metabolic scaling exponents. A three-quarter exponent, observed over many orders of magnitude of body mass, corresponds to a smaller metabolic rate for large animals than proportionality, and to a larger one than surface-limited scaling. Illustrative lines, not measured data.

In 1997 [West, Brown and Enquist](#) derived the three-quarter exponent from three assumptions: that supply networks are space-filling and fractal-like, that terminal units (capillaries) are the same size in all animals, and that the energy needed to distribute resources is minimised. This ties the branching geometry of Section 4.4 to a quantitative law. The model has been criticised: the measured exponent varies between groups, and some analyses find values nearer to two-thirds. A [unifying explanation by Glazier](#) proposes that the exponent shifts with the balance between volume-related and surface-related constraints. Whichever view is correct, the pattern is a power law with an exponent fixed by transport geometry. Evidence grade: **B** for the law; **C** for the universality of the exponent.

5.6 Body plans and morphospace

D'Arcy Thompson's insight was that a form can be treated as a point in a space of parameters. David Raup applied the idea to mollusc shells in 1966. A coiled shell can be described by only a few parameters: the rate W at which the whorl expands, the distance D of the generating curve from the coiling axis and the rate T of translation along the axis. Systematically varying them generates the *theoretical morphospace* of all possible coiled shells.




Figure 20. Raup-style shell coiling. A circular generating curve is expanded at rate W per whorl and offset from the axis by fraction D . Different parameter values produce open, tight and flat coils. Original construction with illustrative parameters.

The result is that real shells occupy only part of the space. [Raup](#) found that real shells occupy only part of the theoretical space and leave large regions empty. Some empty regions plausibly hold forms that would work poorly, but others appear to be viable shapes that no lineage has explored. This is the signature of Layer 2 and 3 constraints working together: some forms are excluded by function, others by the way development and evolution happen to proceed.

A broader example is the *absence* of wheels. Rotating joints exist at the molecular level, in the bacterial flagellar motor, but no animal has a wheel with an axle. The usual explanation is that a freely rotating organ cannot be connected to the rest of the body by blood vessels and nerves, so the form is physically possible but developmentally unreachable. A similar argument is made for the tetrapod body plan, with four limbs and at most five digits in most living species. The earliest limbed vertebrates had more than five digits, so the five-digit pattern is not a physical necessity; it was fixed later in evolution and then conserved. Evidence grade: **B**.

5.7 From gene to geometry: the pinecone revisited

The case of the pinecone now has a full causal chain across three layers. At the biological layer, genes set the production, transport and sensing of auxin and the rate of growth of the shoot tip. At the geometric layer, auxin depletion around each primordium produces an inhibition field. At the dynamical layer, the inhibition field makes the golden angle the attractor of the placement process. At the fundamental layer, the golden angle is the angle that packs a disc uniformly. The Fibonacci spiral counts are the visible result.

Because the final geometry is fixed by the attractor and not by any one gene, many different mutations change the details (cone size, scale shape, spiral handedness) without changing the Fibonacci character. This is why very different plant lineages show the same pattern, and why it reappears in such different organs as pinecones, sunflowers, succulents and cacti.

Key points. Biology does not fight physics; it recruits it. Conserved genes supply parameters, morphogen gradients and reaction–diffusion systems turn them into spatial pattern, and cell mechanics shapes the tissue. The result is a developmental system that produces the same forms from many genetic backgrounds, with large regions of morphospace left unexplored. Constraint from development is real, but it is a bias and not a prohibition.

6. Emergent Structures: Order Without a Designer

Layer 4. Crystals, convection patterns, collective animal behaviour and other complex phenomena show how large-scale organisation can emerge without a central designer specifying every detail.

6.1 What emergence means here

A system is *self-organising* when global order appears from local interactions without any element possessing a plan of the whole. The physicist Philip Anderson summarised the idea in [More Is Different](#): at each level of complexity, new properties appear that cannot be read off from the properties of the parts alone, but are nevertheless fully consistent with them.

Three ingredients appear again and again. The first is an *energy or matter flux*: a system far from equilibrium, which Ilya Prigogine called a [dissipative structure](#). The second is *feedback* (positive feedback amplifies small differences, negative feedback limits them). The third is *local interaction*, in which each element responds only to its neighbours. Given these, patterns with characteristic length scales can emerge and are *selected by the dynamics*, not by any outside specification.

This is the layer at which the claim that a designer need not specify every detail is clearest, because the processes can be simulated, and the simulated forms compared with the natural ones.

6.2 Crystals and snowflakes

A snowflake is the paradigm case of a form that is both highly ordered and unique. Its six-fold symmetry comes from the hexagonal lattice of ice (Section 3.4). Its *shape* depends on the conditions in which it grows. As documented in the [physics of snow crystals](#) by Kenneth Libbrecht, the habit changes with temperature and humidity: thin plates near minus 2 degrees Celsius, hollow columns near minus 5, and thin plates and branched dendrites near minus 15. Higher humidity favours more branching, because a corner sticks out into more vapour and grows faster than a flat face.

Why are the six arms nearly identical? The arms do not communicate. They are exposed to nearly the same conditions as the crystal drifts through the cloud, and they follow the same growth rule, so they develop in parallel. Why are no two snowflakes alike? Because their paths through the cloud differ, and the small differences accumulate. The same logic explains why the two sides of an animal are nearly but not exactly alike.

A cellular model by [Clifford Reiter](#), published in 2005, captures this with a few rules on a hexagonal grid: water vapour diffuses freely; a cell freezes once enough water has accumulated; and vapour is added at a rate γ only at cells that are frozen or touch the crystal, on top of a background vapour level β .




Figure 21. Snow-crystal growth in a Reiter-type cellular model on a hexagonal grid. Only the background vapour level β , the vapour addition γ and the run length change between panels. Higher β gives compact stars; lower β gives more branched dendrites. Original simulation.

The model produces six-fold symmetric shapes from a single seed, plates and dendrites from different parameters, and no two runs with noise would be identical. Evidence grade: **A** for the symmetry origin; **B** for the habit diagram; **B** for the cellular model as a representation.

6.3 Patterns in fluids, chemistry and sand

Pattern formation under flux is widespread, and a few cases show how general the logic is.

Table 7. Pattern formation far from equilibrium

System	Driver	Pattern	What sets the scale
Rayleigh–Bénard convection	Heating a fluid layer from below	Rolls or hexagonal cells	Depth of the fluid layer
Belousov–Zhabotinsky reaction	Chemical energy	Spiral and target waves	Reaction and diffusion rates
Wind ripples and dunes	Wind flow over sand	Ripples, crescent dunes	Distance a grain travels in a hop
Banded vegetation	Water running downslope	Stripes and spots	Water redistribution distance

In each, a uniform state becomes unstable above a threshold of driving, and a pattern with a characteristic wavelength grows. The same equations appear (amplitude equations near threshold), which is why a convection roll, a sand ripple and a stripe of vegetation resemble each other. The semi-arid vegetation case, modelled by [Klausmeier](#), is particularly clear: a simple pair of equations for water and plant biomass reproduces the observed bands and their slow uphill migration.

Evidence grade: **A** for convection and BZ; **B** for vegetation and dunes.

6.4 Collective motion

Flocks of starlings, schools of fish and herds of wildebeest move as if they were one body. [Craig Reynolds](#) showed in 1986 that flock-like motion can be generated by three rules for each individual: avoid collisions, match neighbours' velocity and stay close to the group. A minimal physical version is the model of [Vicsek and colleagues](#) (1995): particles move at constant speed and at each step

adopt the average direction of neighbours within a fixed radius, plus random noise. At low noise the whole group aligns; at high noise it is disordered, with a transition in between.




Figure 22. A Vicsek-type simulation with 300 particles in a periodic box. The order parameter, the length of the mean velocity vector, falls from near one to near zero as noise rises. Snapshots show aligned motion at low noise and random motion at high noise. Original simulation.

In 2008 [Ballerini and colleagues](#) reconstructed the three-dimensional positions of individual birds in flocks of starlings above Rome and found that each bird interacts with a fixed number of neighbours (about six to seven) and not with all neighbours within a fixed distance. This *topological* interaction keeps flocks cohesive when their density varies. The nature of the Vicsek transition in large systems (continuous or discontinuous) has been debated, but the existence of an ordering transition is not. Evidence grade: **A** for the model; **B** for its application to real flocks.

Insect colonies show a related pattern, [stigmergy](#): individuals modify the environment (laying pheromone) and later individuals respond to the modification. Ants find short routes between nest and food because pheromone accumulates faster on shorter paths. The colony solves an optimisation problem without any ant knowing the problem.

6.5 Simple rules, complex outputs

The simplest systems with this property are *cellular automata*, in which a row or grid of cells updates each step according to a fixed rule based on neighbours. In an elementary cellular automaton each cell has two states and sees three cells, so there are only 256 rules. [Stephen Wolfram](#) classified their behaviour in the 1980s.




Figure 23. Three elementary cellular automata. Rule 90 generates the Sierpiński triangle from one cell. Rule 30 generates an irregular pattern that is used as a source of pseudo-random numbers. Rule 110 shows localised moving structures. Original simulation.

[Rule 110](#) was proved by Matthew Cook to be [computationally universal](#), meaning that in principle it can run any computation. The lesson for the thesis is that extremely small rule sets can generate an unbounded range of outputs, and also that outputs of different complexity can arise from rules of the same size. Conway's [Game of Life](#), with two states and one rule, produces moving "gliders", stable "blocks" and many other recurring structures.

The pigmentation of some mollusc shells, such as [Conus textile](#), resembles the output of a one-dimensional automaton. The resemblance is suggestive but resemblance alone is the kind of evidence that Section 1.4 warns against: models for shell pigmentation based on reaction–diffusion also reproduce it. Evidence grade: **A** for the mathematics; **C** for biological interpretation.

6.6 Criticality and power laws

Some systems organise themselves to the edge of a phase transition without being tuned there. In their 1987 model of a [sandpile](#), Per Bak, Chao Tang and Kurt Wiesenfeld dropped grains one at a time onto a grid. Any site with four grains topples, passing one grain to each neighbour, which may trigger further toppling. The result is avalanches of all sizes with a power-law distribution, without any tuning parameter: [self-organised criticality](#).




Figure 24. Avalanche sizes in a simulated Bak–Tang–Wiesenfeld sandpile on a 64 by 64 grid, after dropping 600,000 grains (first 50,000 discarded). Probability density against number of topplings, with log-spaced bins. The fitted slope over the range shown is close to minus one, with a cutoff at the system size. Original simulation.

The idea has been applied to earthquakes, forest fires, neuronal activity and ecosystems. It is also a standing cautionary example. Many reported power laws in empirical data do not survive rigorous testing. [Clauset, Shalizi and Newman](#) showed that distinguishing a true power law from a log-normal or stretched exponential requires careful statistics over a wide range, which is often unavailable. Sandpile models give clean power laws, while real-world cases are mixed. Evidence grade: **A** for the models; **B** for earthquakes (the Gutenberg–Richter law is a well-tested power law); **C** for most other claimed applications.

6.7 Type versus token

Emergent structures sharpen a distinction that the rest of the report relies on: recurrence is of *types*, not *tokens*. No two zebra stripe patterns are the same, yet every zebra shows a pattern of the same kind, with a characteristic width and branching. No two snowflakes are alike, yet every one has six arms. Two Ising lattices will never match spin for spin, yet their cluster statistics are identical.

When asking why forms recur, the right question is therefore about *classes*: why is this class of form reached by so many systems? The answer, at the emergent layer, is that the class is an attractor of a local dynamics, and details are set by fluctuations. This is why the report makes claims about *recurrence of pattern types* and not about the identity of instances.

Key points. Emergent structures are the clearest demonstration that complex order needs no central specification. Flux, feedback and local interaction produce patterns with characteristic scales; the same few mechanisms (convection, reaction–diffusion, alignment, criticality) recur across physics, chemistry and biology. Individual instances differ, but classes of pattern are stable attractors. The principal caution is statistical: resemblance and apparent power laws must be tested, not asserted.

7. Reproductive and Evolutionary Structures: Why Nature Repeats Itself

Layer 5. Shared ancestry, conserved developmental programmes, genetic constraints and convergent evolution explain why organisms can be nearly identical, or independently evolve similar features.

7.1 Heredity as high-fidelity copying

Life adds something that no physical pattern-forming system has: a mechanism that copies the instructions for form with very high fidelity. The copying is what turns a single successful outcome into a lineage, and a lineage into a species. Fidelity is achieved in layers.

The replicating DNA polymerase selects the correct nucleotide with an error rate of roughly one in ten thousand to one in a hundred thousand. A proofreading activity that removes mismatched nucleotides lowers the error rate by two to three orders of magnitude. A post-replication [mismatch repair](#) system lowers it again by a similar factor. [Kunkel's review](#) gives the details. The result is an error rate per base per replication of about one in a billion or lower. The observed [germline mutation rate in humans](#) is higher, at about one per hundred million bases per generation, because it integrates many cell divisions and damage that is not replication error. A child inherits roughly sixty new point mutations among three billion bases.




Figure 25. Approximate error rates in DNA replication at successive levels of correction, and the observed human germline mutation rate. Ranges are order-of-magnitude values from the literature (Kunkel, 2004; Kong et al., 2012). Illustrative compilation.

High-fidelity copying makes the lineage a stable carrier of whatever the generator produced, which allows conserved programmes to persist for hundreds of millions of years. It is the reason that the same Hox genes appear in a mouse and a fly. Evidence grade: **A**.

7.2 Three routes to identical form

The question of why animals "can be identical" has three distinct answers, and the first step in any analysis is to decide which one applies.

Route 1: identical genome. [Monozygotic twins](#) arise when one embryo splits, at a rate of about three to four per thousand births. Clonal organisms are the extreme case. The aspen colony known as [Pando](#) in Utah is a single genetic individual of tens of thousands of stems connected by a common root system. Bacterial colonies, aphid clones and cuttings of plants are further examples.

Even here, identical genomes do not produce identical individuals. Twins differ in fingerprints and in the [epigenetic marks](#) that control gene activity, which diverge as they age. The cause is developmental noise: stochastic events in cell division and signalling that no genome can specify. This is the type-versus-token distinction of Section 6.7 applied to organisms.

Route 2: shared ancestry. Members of a species are similar because they inherit the same regulatory programmes from a recent common ancestor, and selection removes variants that disrupt them. At deeper levels, the forelimbs of a bat, a whale and a human have the same bones in the same arrangement because they are modifications of the same inherited limb programme. This is *homology*.

Route 3: shared constraint and shared selection. Similar forms may appear in lineages with no recent common ancestor because the same problem has the same best solutions. This is *convergence*, and it is the route in which creation structures do the most work.

Table 8. Three routes to similar form

Route	What is shared	Test	Example
Identical genome	Whole genome (copy)	Genetic identity	Twins, Pando, clones
Shared ancestry (homology)	Inherited developmental programme	Phylogeny; conserved genes	Tetrapod limb bones
Convergence	Constraint plus selective problem	Independent origins on a tree	Streamlined bodies; camera eyes

7.3 Deep homology

The three routes interact. In 1997 [Shubin, Tabin and Carroll](#) argued that the genes controlling the development of fish fins and tetrapod limbs are shared, and in 2009 they proposed [deep homology](#): organs that are not homologous as structures can nevertheless be built using homologous genetic regulatory circuits. The paradigm case is the eye. The camera eye of a vertebrate and the compound eye of an insect have independent evolutionary origins, but both depend on Pax6 for their development.

Deep homology complicates the simple split between homology and convergence, and it supports the thesis in two ways. First, it shows that some convergence is not independent at the molecular level: an ancient toolkit is *reused*, so that independent lineages start with the same raw materials. Second, it shows that the generators are deeply shared even when forms diverge, which implies that the biases in what can be produced are shared as well. The [evolution of the eye](#) is a classic instance, with different lineages elaborating light-sensitive cells into more complex organs along repeated paths. Evidence grade: **B**.

7.4 Convergent evolution

Where lineages face the same problem, they often reach the same solution. The best-documented examples are in the following table.

Table 9. Documented convergences

Trait	Independent origins	Source
Camera-type eyes	Vertebrates and cephalopods, among at least 40 independent origins of eyes of some kind	Fernald 2006
Powered flight	Insects, pterosaurs, birds and bats (four origins)	Wikipedia
C4 photosynthesis	More than 60 origins in flowering plants	Sage et al. 2011
Streamlined fusiform body	Ichthyosaurs, dolphins, tuna and sharks	Wikipedia
Crab-like body (carcinisation)	At least five times in decapod crustaceans	Wikipedia
Sabre teeth	Several cat and non-cat lineages, including a marsupial	Wikipedia
Antifreeze glycoproteins	Arctic cod and Antarctic notothenioid fish	Chen et al. 1997
Echolocation	Bats and toothed whales, with convergent changes in the hearing gene prestin	Liu et al. 2010




Figure 26. Convergent body form. Amber tips show the fusiform body. The ichthyosaur and the dolphin each re-evolved it from land-dwelling ancestors; the origin of the shape within fishes depends on ancestral reconstruction. Simplified, schematic phylogeny.

The strength of convergence as evidence depends on the independence of the origins, which is established by mapping the trait on a phylogeny. Two points need care. First, the *molecular* convergence reported for echolocation was originally reported by [Parker and colleagues](#) as widespread across the genome, but [Thomas and Hahn](#) showed that comparable levels of convergence are found in animals that do not echolocate, so the genome-wide signal was largely explained by the statistical null model used. Second, convergence in outward form does not require convergence in the genes that build it. Evidence grade: **A** for convergence of form and function; **C** for convergence at the sequence level.

7.5 Adaptive landscapes and morphospace

Sewall Wright's metaphor of the [fitness landscape](#) captures why convergence happens. Imagine the space of possible forms as a surface, with height giving fitness. Populations climb uphill. If the landscape has a small number of high peaks, lineages starting in different places climb the same peaks.




Figure 27. An illustrative fitness landscape with three major peaks and two minor ones. Lineages starting from different points (white) climb by noisy gradient ascent and finish (amber) at the same few peaks. Illustrative simulation, not empirical data.

The shape of the landscape comes from the lower layers. The positions of the peaks are set by physics: a body that moves through water has an optimal shape fixed by fluid dynamics, a lens has an optimal curvature fixed by optics, and a vessel has an optimal branching fixed by Murray's law. The *accessibility* of the peaks is set by development: only forms that the generator can produce can be tested by selection. If the landscape is rugged, with many peaks, different lineages may find different ones, and history matters. If it is smooth with few peaks, convergence is the norm.

7.6 Genotype–phenotype maps

A final constraint operates between genes and forms. The mapping from genotype to phenotype is *many-to-one* and highly uneven. In a classic computational study of RNA, [Schuster and colleagues](#) showed that a small number of RNA folds account for most of the possible sequences, while the great majority of shapes are reachable from very few sequences. [Dingle, Camargo and Louis](#) showed that the same strong bias towards simple outputs appears in many input–output maps, including those used in biological modelling.




Figure 28. A schematic genotype–phenotype map. Left: regions of genotype space that map to the same phenotype (colours); a few phenotypes cover large regions. Right: share of genotype space by phenotype, ranked. Illustrative simulation based on weighted Voronoi regions, not empirical data.

The consequence is that variation is not isotropic. The mutations that occur are far more likely to produce a few common phenotypes than rare ones, so evolution tends to arrive repeatedly at the *same* phenotypes. This *developmental bias* adds a second source of recurrence to selection. It also gives a response to the old question of the arrival of the fittest: a selective filter needs variants to choose among, and the map decides which variants are offered. Evidence grade: **B**.

7.7 Contingency and convergence

How strong is convergence relative to contingency? Stephen Jay Gould argued in [Wonderful Life](#) that if the tape of life were replayed, the outcome would differ, because small accidents (an extinction, a mutation) change what follows. [Simon Conway Morris](#) argued in *Life's Solution* that convergence is so pervasive that replaying the tape would produce similar kinds of forms, if not the same species.

The most direct experiment is the [Long-Term Evolution Experiment](#), begun by Richard Lenski in 1988, in which twelve populations of *Escherichia coli* started from one clone and have evolved separately for tens of thousands of generations. In the first ten thousand generations, [fitness rose in parallel](#) in all twelve populations, while cell size and other traits diverged. Later, one population evolved the ability to use citrate, a rare event that [depended on earlier chance mutations](#) and did not occur in the other eleven.

The result supports both authors at different levels. Convergence is strong for *function and the class of solution* (high fitness in a given environment), where the possibility space is narrow. Contingency is strong for *lineage-specific details*, where the space is wide and the landscape is rugged. This division is the heart of the creation-structure view of evolution: history chooses among a restricted menu. Evidence grade: **B**.

Key points. Heredity turns outputs of generators into persistent lineages. Similar forms arise by three routes (shared genome, shared ancestry, shared constraint and selection), and the routes can be told apart on a phylogeny. Convergence is common because fitness landscapes have few high peaks, and the genotype–phenotype map biases variation towards the same few outcomes. Contingency is real but operates within these limits.

8. Synthesis: A Unified Framework

8.1 One architecture, five layers

The previous five sections described creation structures one layer at a time. This section combines them. The core proposition is that a recurring form is the output of a pipeline with four stages.

1. **Constraint.** Mathematics and physics define the space of possible forms (Layers 1 and 2).
2. **Generation.** A developmental or dynamical process explores that space with a bias, producing forms from parameters and noise (Layers 3 and 4).
3. **Heredity.** Parameters of the generator are copied with high fidelity, so a successful configuration can persist (Layer 5).
4. **Selection.** The environment filters the outputs, leaving those that perform (Layer 5).

Proposition (grade D). The probability that independent systems produce the same form class is higher when (i) the viable region of form space is small, (ii) the generator has a large basin of attraction for that class, (iii) the selective landscape has few peaks, and (iv) heredity preserves the generator's parameters faithfully.

This is a qualitative framework, not a formula, and it is offered as a hypothesis whose individual terms can be measured (Section 9). It has the advantage that each of the four conditions maps onto a body of existing methods: the size of the viable region can be estimated by sampling and simulation, basins by perturbation analysis, peaks by experimental or computational landscapes, and fidelity by sequencing.

8.2 Case studies across the layers

The figure below applies the framework to five recurring forms and to identical twins. For each, it shows the relative contribution of each layer, assessed from the literature discussed in this report.

	Pinecone spirals	Honeycomb	Snowflake	Zebrafish stripes	Convergent body shape	Identical twins
Fundamental (law, symmetry)	moderate	dominant	dominant	minor	dominant	minor
Geometric (packing, growth)	dominant	dominant	moderate	minor	minor	–
Biological (gene, development)	dominant	minor	–	dominant	moderate	dominant
Emergent (self-organisation)	moderate	moderate	dominant	dominant	–	–
Evolutionary (heredity, selection)	moderate	moderate	–	moderate	dominant	dominant

Figure 29. Relative contribution of each layer to five recurring phenomena and to the case of identical twins. The ratings (dominant, moderate, minor, none) are the author's qualitative judgement from the evidence reviewed in Sections 3 to 7, and not measured quantities.

Pinecone spirals are driven by geometry (Layer 2) and developmental dynamics (Layer 3): the golden angle is the unique solution to the packing problem and the attractor of the inhibitory-field dynamics. Selection and heredity matter because they maintain the machinery, but they do not need to select the pattern itself.

Honeycomb is dominated by constraint (Layers 1 and 2). The hexagon is optimal and also the natural result of surface tension on wax. Behaviour (Layer 5, as bees' building instincts evolved) supplies the process, but the form is dictated by geometry.

Snowflakes are the most purely physical case: the lattice (Layer 1), the shape of the growing face and diffusion-limited growth (Layers 2 and 4). Heredity and selection play no role at all, which is why snowflakes show that recurrence needs neither.

Zebrafish stripes and digit patterns are developmental, with reaction–diffusion dynamics (Layers 3 and 4) as the generator and a heavy contribution from heredity (the parameters are tuned genetically, as shown by the Hox dose experiments).

Convergent body shape depends on physics (the hydrodynamic optimum) and on selection, with development providing the route.

Identical twins are special: nearly everything is carried by heredity (a copy of the genome) and development (running the same programme twice), and the divergence between twins arises from noise.

8.3 Interactions between layers

The layers are not independent. Four interactions deserve emphasis.

- **Constraint shapes the generator.** Because physics fixes what works, a generator that produces non-viable forms is removed. Over time, generators are those compatible with physics.

- **Evolution tunes physical generators.** The Hox-dose experiment shows evolution acting by changing the *parameter* of a physical process (the wavelength of a Turing pattern) and thus moving along a continuous family of forms. This is a far smaller step than building a new structure from scratch, which explains why evolution is fast when it follows an existing generator.
- **Selection acts through the landscape that physics defines.** The shape of the hydrodynamic optimum is set by fluid dynamics, so selection can only follow it.
- **Noise creates tokens from types.** Fluctuations in development create individual differences, leaving the class of form intact.

8.4 Why so much order?

The observation that the natural world is surprisingly ordered, which motivated this report, can now be answered in three parts.

1. **Few ingredients.** Identical particles with fixed interactions mean that the same building blocks recur everywhere.
2. **Few attractors.** Many different processes, including packing, inhibition, diffusion with feedback and alignment, have a small number of stable outcomes, so that many starting points converge.
3. **Few solutions.** Selection and optimisation, applied to problems with narrow solutions (move through water, collect light, deliver blood), keep only a few of the forms the generators can produce.

Each answer is checkable, and each fails somewhere: some patterns are chance (Section 9.3), and some similarities are due to shared history. But the three together account for much of what is observed.

Key points. A recurring form is the product of a pipeline: constraint, generation, heredity and selection. The layers contribute unequally to each phenomenon, and the contribution can be assessed by asking what would happen if each layer were removed. Evolution typically works by tuning parameters of generators that physics already supplies.

9. Testing the Thesis: Predictions, Limits and Risks

A thesis that cannot fail is not a scientific thesis. This section states what the creation-structure view predicts, what would count against it and where it is most at risk of being wrong.

9.1 Testable predictions

Table 10. Eight predictions of the creation-structure thesis

#	Prediction	How to test	Status
P1	Lineages that independently solve the same problem show form similarity beyond what chance and shared ancestry predict.	Morphometrics across independent origins on a phylogeny.	Supported in cases (Section 7.4); no systematic survey.
P2	A minimal generative model reproduces a recurring form class without form-specific tuning.	Simulation (e.g. Vogel, Gray–Scott, Reiter models).	Supported for phyllotaxis, snow crystals, stripes.
P3	Changing a constraint parameter shifts the form in the predicted direction.	Perturbation (Hox dose and digit number; droplet timing and divergence angle).	Supported.
P4	Regions of form space excluded by physics remain empty; regions excluded only by development can be populated by engineering.	Synthetic morphogenesis; organoids; shell and venation morphospaces.	Early evidence.
P5	Genotype–phenotype maps are strongly biased towards a few phenotypes.	Computational maps; deep mutational scans.	Supported (RNA, model systems).
P6	The number of independent origins of a trait correlates with the size of its basin of attraction.	Compare origin counts with simulated basin sizes.	Untested systematically.
P7	Where selection is weak, forms diverge more, beyond the limits set by constraint.	Compare constrained and unconstrained traits across clades.	Partly supported.
P8	A pattern fails to appear in lineages lacking the generating mechanism.	Survey Fibonacci phyllotaxis against meristem inhibition mechanisms.	Untested.

9.2 What would count against the thesis

Three kinds of finding would weaken the framework substantially.

- 1. Recurrence without mechanism.** If a recurring pattern is robustly found but no generator can reproduce it, the framework would have no explanation to offer, and an explanation in terms of historical accident would gain weight.
- 2. Constraint-free production.** If a system lacking the supposed constraint reliably produces the same form (for example, hexagonal order arising in a system without any packing or tension effect), the constraint is not the cause.

3. **Easy escape.** If engineers can readily and stably produce forms in regions of morphospace that the framework treats as inaccessible, using the same developmental system, then the constraint is weaker than claimed.

9.3 Failure modes

The history of the subject contains many examples of explanations that were too good to be true. Six failure modes should be guarded against.

Golden-ratio inflation. Claims that the golden ratio governs the proportions of the Parthenon, the human face or the stock market are mostly false or unfalsifiable. George Markowsky's [survey of misconceptions](#) documents many, including mismeasured buildings and selective choice of landmarks. The golden ratio matters in phyllotaxis because a mechanism connects it to the observation, and not otherwise.

Pareidolia and selective sampling. The human visual system finds familiar shapes in noise. Collections of examples chosen because they fit are not evidence of a law. Always ask what the *denominator* is: how many systems were examined and how many showed the pattern?

Adaptationist storytelling. [Gould and Lewontin](#) criticised the habit of explaining every trait as a direct adaptation. Features may be by-products of constraint, like the architectural *spandrels* of a cathedral, which are unavoidable consequences of arches. The creation-structure view *agrees* with this critique and gives it a constructive form: ask first what the constraints force.

Power-law and fractal overreach. As discussed in Sections 4.5 and 6.6, claims of scale-free behaviour need a wide scaling range and rigorous statistics.

Scale confusion. The same geometry at different scales (hurricane and galaxy spirals) may have unrelated causes. Resemblance across scales is a hypothesis, not a result.

Contingency denial. The framework should not be used to claim that everything was inevitable. The citrate experiment of Section 7.7 shows that contingent events happen even under tightly controlled conditions.

9.4 Open questions

Five questions will decide how far the framework can be developed.

1. **Measuring possibility spaces.** How can the size of the viable region of form space be estimated for real morphologies, such as leaf shapes, wing venation or skeletons?
2. **Quantifying generator bias.** Can the strength of developmental bias be measured and compared across lineages, in units that allow a prediction of recurrence rates?
3. **Linking genes to attractors.** How do changes in gene networks map onto changes in the parameters of physical generators, such as the wavelength of a Turing pattern?
4. **Classifying attractors.** Is there a short list of morphogenetic attractor types, as there is for phase transitions, that covers most recurrent biological patterns?
5. **Evolvability.** Do lineages differ in how easily they explore form space, and is that difference itself heritable and selectable?

9.5 A note on interpretation

The recurrence of mathematical structure in nature has long invited philosophical comment. Eugene Wigner's essay on the [unreasonable effectiveness of mathematics](#) is the best-known statement of the puzzle. Positions range from Platonism (mathematical structures exist independently and nature instantiates them) to structural realism (what physics describes is structure) to the view that mathematics is a human tool that nature happens to suit.

The word *creation* in this report's title is used in the sense of *generation*: the production of form by processes. Nothing in the argument requires or excludes a view about the ultimate origin of the laws themselves, and the scientific content stands on the evidence for mechanisms alone. This is deliberate. The question of why the laws are what they are lies outside the scope of a research programme that must be testable.

10. Applications and Heuristics

10.1 Where creation structures create value

If recurrence reflects a small number of generators, then knowing the generators makes it possible to *use* them. Several fields already do.

Table 11. Applications of creation structures

Field	Principle exploited	Example	Status
Architecture and materials	Soap-film minimisation; hexagonal and Voronoi lattices	Tensile roofs designed with soap-film models; honeycomb sandwich panels	Established
Fluid and transport networks	Murray's law; fractal-like branching	Biomimetic microfluidic and cooling networks	Developing
Computing and AI	Cellular automata; swarm dynamics	Neural cellular automata that grow and repair images; ant colony optimisation	Developing
Synthetic biology	Engineered reaction–diffusion	Stochastic Turing patterns built into bacterial populations	Early
Tissue engineering	Self-organisation of cells	Brain organoids formed from stem cells	Early
Design and biomimetics	Transferring natural solutions	Biomimetics generally	Mixed; claims vary in quality

The creation-structure view suggests a more principled approach to biomimetics than copying shapes. Rather than imitating a form (for example, a leaf), the engineer identifies the *generator* and the *constraint* that produced it, and then re-implements them in a new medium. The distinction matters because a copied shape carries the constraints of its original medium, while a copied generator can be adapted to a new one.

10.2 Ten heuristics for reading a pattern

The following heuristics condense the report into rules that analysts, designers and researchers can apply when they meet a recurring pattern. They are intended to be used in order, but each can stand alone.

Table 12. Ten heuristics

#	Heuristic	Do this	Typical error avoided
H1	Name the layer.	Decide whether the pattern is best explained by law, geometry, development, self-organisation or evolution.	Explaining physics with biology, or the reverse.
H2	Demand a mechanism.	Require a process that generates the pattern, not only a description of it.	Mistaking resemblance for explanation.
H3	Build the null.	Ask what random or unconstrained processes would produce.	Seeing a pattern in noise.
H4	Count independent origins.	Map the trait on a phylogeny or across unrelated systems.	Treating shared ancestry as convergence.
H5	Find the attractor.	Test robustness: does the form return after perturbation?	Confusing a fragile result with a stable one.
H6	Simulate the minimum.	Build the simplest generator that reproduces the class of form.	Over-complicated explanations.
H7	Separate type from token.	Compare classes of form, not individual instances.	Expecting identical instances from identical rules.
H8	Check the range.	Require a wide scaling range before claiming a fractal or power law.	Spurious scale-free claims.
H9	Look at what is missing.	Map the forms that do <i>not</i> occur, and ask why.	Ignoring absent forms that reveal constraints.
H10	Leave room for history.	If no constraint explains the pattern, say so and treat it as contingent.	Forcing inevitability on accidents.




Figure 30. A decision procedure for classifying the likely source of a recurring pattern. The route is a first-pass triage and not a proof; the layers are those defined in Section 2.3.

10.3 Research agenda

The thesis points to a programme of work that can be organised in three horizons.

Table 13. A research programme in three horizons

Horizon	Work	Outputs
Near term (0–12 months)	Compile a catalogue of recurring forms with the independent-origin count, mechanism, layer and evidence grade for each.	An open dataset; a reproducible scoring scheme.
Medium term (1–3 years)	Measure possibility-space size and generator bias for tractable cases (shell coiling, wing venation, leaf margins); run controlled perturbation studies.	Quantitative tests of P1, P3, P5 and P6.
Longer term (3–5 years)	Engineer forms in predicted-inaccessible and predicted-accessible regions of morphospace using synthetic or organoid systems.	A direct test of P4; design rules for synthetic morphogenesis.

11. Conclusion

The question that began this report was why nature repeats itself. The answer proposed is that repetition is the visible trace of a layered architecture. Mathematics and physics restrict the forms that can exist. Generative processes, including packing, diffusion with feedback, alignment, development and growth, explore the permitted forms with a strong bias, so that a handful of outcomes are reached again and again from different starting points. Heredity copies the parameters of successful generators with extraordinary fidelity, and selection filters what they produce.

This account is pluralist. The spirals of a pinecone, the hexagons of a honeycomb, the arms of a snowflake and the body of a dolphin have different causes. What they share is a *structure of explanation*: each is the product of a small possibility space, a biased generator and, in the living cases, a filter. The five-layer model gives that structure a vocabulary, and the ten heuristics give it a practice.

The account is also testable and limited. It predicts that independent systems solving the same problem will converge, that minimal generative models will reproduce recurring forms, that perturbing constraints will shift forms predictably and that genotype–phenotype maps will be biased. Many of these predictions are already supported, and a few have yet to be tested. It does not claim that everything is inevitable: contingency operates within the constraints, as the citrate experiment shows.

The practical consequence is a change in the question asked of any pattern. Instead of asking only *what does this resemble*, the creation-structure view asks *which constraints, generator and filter produced this, and would a different system with the same constraints produce it too?* A pattern for which the answer is known can be predicted, simulated and, increasingly, engineered.

References

Primary literature

The works are listed alphabetically by first author. Each entry links to the publisher's record through its DOI or to the open preprint.

- Anderson, P. W. (1972). More is different. *Science* 177, 393–396. [doi:10.1126/science.177.4047.393](https://doi.org/10.1126/science.177.4047.393)
- Avnir, D., Biham, O., Lidar, D., Malcai, O. (1998). Is the geometry of nature fractal? *Science* 279, 39–40. [doi:10.1126/science.279.5347.39](https://doi.org/10.1126/science.279.5347.39)
- Bak, P., Tang, C., Wiesenfeld, K. (1987). Self-organized criticality: an explanation of 1/f noise. *Physical Review Letters* 59, 381–384. [doi:10.1103/PhysRevLett.59.381](https://doi.org/10.1103/PhysRevLett.59.381)
- Ballerini, M., et al. (2008). Interaction ruling animal collective behavior depends on topological rather than metric distance. *PNAS* 105, 1232–1237. [doi:10.1073/pnas.0711437105](https://doi.org/10.1073/pnas.0711437105)
- Blount, Z. D., Borland, C. Z., Lenski, R. E. (2008). Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *PNAS* 105, 7899–7906. [doi:10.1073/pnas.0803151105](https://doi.org/10.1073/pnas.0803151105)
- Chen, L., DeVries, A. L., Cheng, C.-H. C. (1997). Convergent evolution of antifreeze glycoproteins in Antarctic notothenioid fish and Arctic cod. *PNAS* 94, 3817–3822. [doi:10.1073/pnas.94.8.3817](https://doi.org/10.1073/pnas.94.8.3817)
- Clauset, A., Shalizi, C. R., Newman, M. E. J. (2009). Power-law distributions in empirical data. *SIAM Review* 51, 661–703. [doi:10.1137/070710111](https://doi.org/10.1137/070710111)
- Cook, M. (2004). Universality in elementary cellular automata. *Complex Systems* 15, 1–40. [PDF](#)
- Davidson, E. H., Erwin, D. H. (2006). Gene regulatory networks and the evolution of animal body plans. *Science* 311, 796–800. [doi:10.1126/science.1113832](https://doi.org/10.1126/science.1113832)
- Dingle, K., Camargo, C. Q., Louis, A. A. (2018). Input–output maps are strongly biased towards simple outputs. *Nature Communications* 9, 761. [doi:10.1038/s41467-018-03101-6](https://doi.org/10.1038/s41467-018-03101-6)
- Douady, S., Couder, Y. (1992). Phyllotaxis as a physical self-organized growth process. *Physical Review Letters* 68, 2098–2101. [doi:10.1103/PhysRevLett.68.2098](https://doi.org/10.1103/PhysRevLett.68.2098)
- Driever, W., Nüsslein-Volhard, C. (1988). A gradient of bicoid protein in *Drosophila* embryos. *Cell* 54, 83–93. [doi:10.1016/0092-8674\(88\)90182-1](https://doi.org/10.1016/0092-8674(88)90182-1)
- Fernald, R. D. (2006). Casting a genetic light on the evolution of eyes. *Science* 313, 1914–1918. [doi:10.1126/science.1127889](https://doi.org/10.1126/science.1127889)
- Fraga, M. F., et al. (2005). Epigenetic differences arise during the lifetime of monozygotic twins. *PNAS* 102, 10604–10609. [doi:10.1073/pnas.0500398102](https://doi.org/10.1073/pnas.0500398102)
- Gibson, M. C., Patel, A. B., Nagpal, R., Perrimon, N. (2006). The emergence of geometric order in proliferating metazoan epithelia. *Nature* 442, 1038–1041. [doi:10.1038/nature05014](https://doi.org/10.1038/nature05014)
- Gierer, A., Meinhardt, H. (1972). A theory of biological pattern formation. *Kybernetik* 12, 30–39. [doi:10.1007/BF00289234](https://doi.org/10.1007/BF00289234)
- Glazier, D. S. (2010). A unifying explanation for diverse metabolic scaling in animals and plants. *Biological Reviews* 85, 111–138. [doi:10.1111/j.1469-185X.2009.00095.x](https://doi.org/10.1111/j.1469-185X.2009.00095.x)
- Gould, S. J., Lewontin, R. C. (1979). The spandrels of San Marco and the Panglossian paradigm. *Proceedings of the Royal Society B* 205, 581–598. [doi:10.1098/rspb.1979.0086](https://doi.org/10.1098/rspb.1979.0086)
- Halder, G., Callaerts, P., Gehring, W. J. (1995). Induction of ectopic eyes by targeted expression of the *eyeless* gene in *Drosophila*. *Science* 267, 1788–1792. [doi:10.1126/science.7892602](https://doi.org/10.1126/science.7892602)
- Hales, T. C. (2001). The honeycomb conjecture. *Discrete & Computational Geometry* 25, 1–22. [arXiv:math/9906042](https://arxiv.org/abs/math/9906042)
- Hales, T., et al. (2017). A formal proof of the Kepler conjecture. *Forum of Mathematics, Pi* 5, e2. [doi:10.1017/fmp.2017.1](https://doi.org/10.1017/fmp.2017.1)
- Karig, D., et al. (2018). Stochastic Turing patterns in a synthetic bacterial population. *PNAS* 115, 6572–6577. [doi:10.1073/pnas.1720770115](https://doi.org/10.1073/pnas.1720770115)
- Karihaloo, B. L., Zhang, K., Wang, J. (2013). Honeybee combs: how the circular cells transform into rounded hexagons. *Journal of the Royal Society Interface* 10, 20130299. [doi:10.1098/rsif.2013.0299](https://doi.org/10.1098/rsif.2013.0299)

- Klausmeier, C. A. (1999). Regular and irregular patterns in semiarid vegetation. *Science* 284, 1826–1828. [doi:10.1126/science.284.5421.1826](https://doi.org/10.1126/science.284.5421.1826)
- Kleiber, M. (1932). Body size and metabolism. *Hilgardia* 6, 315–353. [doi:10.3733/hilg.v06n11p315](https://doi.org/10.3733/hilg.v06n11p315)
- Kondo, S., Asai, R. (1995). A reaction–diffusion wave on the skin of the marine angelfish *Pomacanthus*. *Nature* 376, 765–768. [doi:10.1038/376765a0](https://doi.org/10.1038/376765a0)
- Kondo, S., Miura, T. (2010). Reaction-diffusion model as a framework for understanding biological pattern formation. *Science* 329, 1616–1620. [doi:10.1126/science.1179047](https://doi.org/10.1126/science.1179047)
- Kong, A., et al. (2012). Rate of de novo mutations and the importance of father's age to disease risk. *Nature* 488, 471–475. [doi:10.1038/nature11396](https://doi.org/10.1038/nature11396)
- Kunkel, T. A. (2004). DNA replication fidelity. *Journal of Biological Chemistry* 279, 16895–16898. [doi:10.1074/jbc.R400006200](https://doi.org/10.1074/jbc.R400006200)
- Lancaster, M. A., et al. (2013). Cerebral organoids model human brain development and microcephaly. *Nature* 501, 373–379. [doi:10.1038/nature12517](https://doi.org/10.1038/nature12517)
- Lenski, R. E., Travisano, M. (1994). Dynamics of adaptation and diversification: a 10,000-generation experiment with bacterial populations. *PNAS* 91, 6808–6814. [doi:10.1073/pnas.91.15.6808](https://doi.org/10.1073/pnas.91.15.6808)
- Libbrecht, K. G. (2005). The physics of snow crystals. *Reports on Progress in Physics* 68, 855–895. [doi:10.1088/0034-4885/68/4/R03](https://doi.org/10.1088/0034-4885/68/4/R03)
- Liu, Y., et al. (2010). Convergent sequence evolution between echolocating bats and dolphins. *Current Biology* 20, R53–R54. [doi:10.1016/j.cub.2009.11.058](https://doi.org/10.1016/j.cub.2009.11.058)
- Mandelbrot, B. (1967). How long is the coast of Britain? Statistical self-similarity and fractional dimension. *Science* 156, 636–638. [doi:10.1126/science.156.3775.636](https://doi.org/10.1126/science.156.3775.636)
- Markowsky, G. (1992). Misconceptions about the golden ratio. *College Mathematics Journal* 23, 2–19. [doi:10.2307/2686193](https://doi.org/10.2307/2686193)
- Maynard Smith, J., et al. (1985). Developmental constraints and evolution. *Quarterly Review of Biology* 60, 265–287. [doi:10.1086/414425](https://doi.org/10.1086/414425)
- Mordvintsev, A., Randazzo, E., Niklasson, E., Levin, M. (2020). Growing neural cellular automata. *Distill.* distill.pub/2020/growing-ca
- Murray, C. D. (1926). The physiological principle of minimum work. I. The vascular system and the cost of blood volume. *PNAS* 12, 207–214. [doi:10.1073/pnas.12.3.207](https://doi.org/10.1073/pnas.12.3.207)
- Nakamasu, A., Takahashi, G., Kanbe, A., Kondo, S. (2009). Interactions between zebrafish pigment cells responsible for the generation of Turing patterns. *PNAS* 106, 8429–8434. [doi:10.1073/pnas.0808622106](https://doi.org/10.1073/pnas.0808622106)
- Parker, J., et al. (2013). Genome-wide signatures of convergent evolution in echolocating mammals. *Nature* 502, 228–231. [doi:10.1038/nature12511](https://doi.org/10.1038/nature12511)
- Pearson, J. E. (1993). Complex patterns in a simple system. *Science* 261, 189–192. [doi:10.1126/science.261.5118.189](https://doi.org/10.1126/science.261.5118.189)
- Raspopovic, J., Marcon, L., Russo, L., Sharpe, J. (2014). Digit patterning is controlled by a Bmp-Sox9-Wnt Turing network modulated by morphogen gradients. *Science* 345, 566–570. [doi:10.1126/science.1252960](https://doi.org/10.1126/science.1252960)
- Reinhardt, D., et al. (2003). Regulation of phyllotaxis by polar auxin transport. *Nature* 426, 255–260. [doi:10.1038/nature02081](https://doi.org/10.1038/nature02081)
- Reiter, C. A. (2005). A local cellular model for snow crystal growth. *Chaos, Solitons & Fractals* 23, 1111–1119. [doi:10.1016/j.chaos.2004.06.071](https://doi.org/10.1016/j.chaos.2004.06.071)
- Sage, R. F., Christin, P.-A., Edwards, E. J. (2011). The C4 plant lineages of planet Earth. *Journal of Experimental Botany* 62, 3155–3169. [doi:10.1093/jxb/err048](https://doi.org/10.1093/jxb/err048)
- Schuster, P., Fontana, W., Stadler, P. F., Hofacker, I. L. (1994). From sequences to shapes and back: a case study in RNA secondary structures. *Proceedings of the Royal Society B* 255, 279–284. [doi:10.1098/rspb.1994.0040](https://doi.org/10.1098/rspb.1994.0040)
- Shechtman, D., Blech, I., Gratias, D., Cahn, J. W. (1984). Metallic phase with long-range orientational order and no translational symmetry. *Physical Review Letters* 53, 1951–1953. [doi:10.1103/PhysRevLett.53.1951](https://doi.org/10.1103/PhysRevLett.53.1951)
- Sheth, R., et al. (2012). Hox genes regulate digit patterning by controlling the wavelength of a Turing-type mechanism. *Science* 338, 1476–1480. [doi:10.1126/science.1226804](https://doi.org/10.1126/science.1226804)
- Shubin, N., Tabin, C., Carroll, S. (1997). Fossils, genes and the evolution of animal limbs. *Nature* 388, 639–648. [doi:10.1038/41710](https://doi.org/10.1038/41710)
- Shubin, N., Tabin, C., Carroll, S. (2009). Deep homology and the origins of evolutionary novelty. *Nature* 457, 818–823. [doi:10.1038/nature07891](https://doi.org/10.1038/nature07891)

- Sick, S., Reinker, S., Timmer, J., Schlake, T. (2006). WNT and DKK determine hair follicle spacing through a reaction-diffusion mechanism. *Science* 314, 1447–1450. [doi:10.1126/science.1130088](https://doi.org/10.1126/science.1130088)
- Steinberg, M. S. (1963). Reconstruction of tissues by dissociated cells. *Science* 141, 401–408. [doi:10.1126/science.141.3579.401](https://doi.org/10.1126/science.141.3579.401)
- Tallinen, T., et al. (2016). On the growth and form of cortical convolutions. *Nature Physics* 12, 588–593. [doi:10.1038/nphys3632](https://doi.org/10.1038/nphys3632)
- Thomas, G. W. C., Hahn, M. W. (2015). Determining the null model for detecting adaptive convergence from genomic data: a case study using echolocating mammals. *Molecular Biology and Evolution* 32, 1232–1236. [doi:10.1093/molbev/msv013](https://doi.org/10.1093/molbev/msv013)
- Turing, A. M. (1952). The chemical basis of morphogenesis. *Philosophical Transactions of the Royal Society B* 237, 37–72. [doi:10.1098/rstb.1952.0012](https://doi.org/10.1098/rstb.1952.0012)
- Vicsek, T., Czirók, A., Ben-Jacob, E., Cohen, I., Shochet, O. (1995). Novel type of phase transition in a system of self-driven particles. *Physical Review Letters* 75, 1226–1229. [doi:10.1103/PhysRevLett.75.1226](https://doi.org/10.1103/PhysRevLett.75.1226)
- West, G. B., Brown, J. H., Enquist, B. J. (1997). A general model for the origin of allometric scaling laws in biology. *Science* 276, 122–126. [doi:10.1126/science.276.5309.122](https://doi.org/10.1126/science.276.5309.122)
- Wigner, E. P. (1960). The unreasonable effectiveness of mathematics in the natural sciences. *Communications on Pure and Applied Mathematics* 13, 1–14. [doi:10.1002/cpa.3160130102](https://doi.org/10.1002/cpa.3160130102)
- Wolfram, S. (1983). Statistical mechanics of cellular automata. *Reviews of Modern Physics* 55, 601–644. [doi:10.1103/RevModPhys.55.601](https://doi.org/10.1103/RevModPhys.55.601)
- Wolpert, L. (1969). Positional information and the spatial pattern of cellular differentiation. *Journal of Theoretical Biology* 25, 1–47. [doi:10.1016/S0022-5193\(69\)80016-0](https://doi.org/10.1016/S0022-5193(69)80016-0)

Books

- Carroll, S. B. (2005). *Endless Forms Most Beautiful: The New Science of Evo Devo*. W. W. Norton. [Reference article](#)
- Conway Morris, S. (2003). *Life's Solution: Inevitable Humans in a Lonely Universe*. Cambridge University Press. [Author page](#)
- Gould, S. J. (1989). *Wonderful Life: The Burgess Shale and the Nature of History*. W. W. Norton. [Reference article](#)
- Mandelbrot, B. B. (1982). *The Fractal Geometry of Nature*. W. H. Freeman. [Reference article](#)
- Thompson, D. W. (1917). *On Growth and Form*. Cambridge University Press. [Full text at the Internet Archive](#)
- Wolfram, S. (2002). *A New Kind of Science*. Wolfram Media. [Free online edition](#)

Glossary

Table 14. Terms used in this report

Term	Meaning
Allometry	The study of how traits scale with body size.
Attractor	A state or set of states towards which a dynamical system evolves from many starting points.
Convergent evolution	Independent evolution of similar traits in lineages without a recent common ancestor.
Creation structure	The rules, constraints and generative processes that determine which forms can emerge, how they develop and why patterns recur.
Deep homology	Shared genetic regulatory circuits underlying structures that are not homologous as organs.
Developmental bias	A tendency of development to produce some variants more readily than others.
Dissipative structure	An ordered structure maintained by a flow of energy or matter through an open system.
Fitness landscape	A surface in form or genotype space whose height gives fitness.
Generator	A process that produces form from parameters and noise.
Genotype–phenotype map	The relationship between a genetic sequence and the form it produces.
Homology	Similarity due to inheritance from a common ancestor.
Morphogen	A signalling molecule whose concentration specifies cell fate by position.
Morphospace	A space whose axes are parameters of form; each point is a possible morphology.
Parastichy	A visible spiral line of neighbouring organs in a phyllotactic pattern.
Phyllotaxis	The arrangement of leaves, seeds or scales on a plant axis.
Reaction–diffusion	A system of interacting, diffusing substances that can generate spatial patterns.
Self-organisation	Emergence of global order from local interactions without a central plan.
Symmetry breaking	The adoption by a system of a state with less symmetry than the laws that govern it.
Universality	Independence of large-scale behaviour from microscopic detail near a phase transition.

Appendix: Key Quantities

Table 15. Quantities cited in this report

Quantity	Value	Where used
Golden ratio, φ	$(1 + \sqrt{5}) / 2 \approx 1.6180$	Sections 3.4, 4.1, 4.2
Golden angle	$360^\circ / \varphi^2 \approx 137.508^\circ$	Section 4.2
Crystallographic rotation orders	2, 3, 4, 6	Section 3.4
Wallpaper, point and space groups	17; 32; 230	Section 3.4
Perimeter of unit-area cell: triangle, square, hexagon	4.56; 4.00; 3.72	Section 4.3
Densest sphere packing	$\pi / \sqrt{18} \approx 74.05\%$	Section 4.3
Murray daughter-to-parent diameter ratio (symmetric split)	$2^{-1/3} \approx 0.794$	Section 4.4
Ising critical temperature (2D, units of J/k)	$2 / \ln(1 + \sqrt{2}) \approx 2.269$	Section 3.5
Kleiber exponent	$\approx 3/4$	Section 5.5
Plateau vertex angle: three films; four edges	120° ; $\approx 109.5^\circ$	Section 3.3
Replication error after proofreading and repair	$\approx 10^{-9}$ to 10^{-10} per base	Section 7.1
Human germline mutation rate	$\approx 1.2 \times 10^{-8}$ per base per generation	Section 7.1
Independent origins of C4 photosynthesis	> 60	Section 7.4